



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

be interesting to know to what extent. Is it a common occurrence, or is it only a rare one? It would also be interesting to know, if any other mosses are likewise thus attacked. That moss spores should be an especial dainty to these little insects, is not at all improbable. Baltimore, Md.

A RED *ANDREÆA*.

CORA H. CLARKE.

On July 3rd, 1906, I was climbing a rocky path in the woods at Manchester, Mass., not far from the ocean, nor much above sea-level, when my eye was caught by the very dark red color of some small cushions of moss on a bit of outcropping rock. At least, they appeared red, or a dark red-brown, by contrast with the other mosses on the same rock. The largest cushion was but little over an inch across. Some of the cushions bore a little fruit, which I did not at first recognize to be that of *Andreæa*; there were spores in some of the capsules.

On July 21st, a visit to this same rock showed the *Andreæa* in a dry condition, and therefore almost black. Not far off was an almost vertical rock-wall, of considerable extent, but with growths of moss here and there. Here again was *Andreæa* but not so much in cushions as in patches with a flat surface. The color was not red at all but black when dry—dark green when wet—with little fruit.

Both forms answer to the description of *Andreæa rupestris*, a species not new to Massachusetts since Mr. Edwin Faxon found it in Franklin Park, Boston, Nov. 25th, 1886. His specimens were red rather than dark green, and showed signs of having borne abundant fruit earlier in the season.

Boston, Mass.

SULLIVANT MOSS CHAPTER NOTES.

Dr. A. J. Grout, 360 Lenox Road, Brooklyn, N. Y., would like to arrange for a Sullivant Moss Chapter exploration of Mt. Washington, N. H., some time during next July. Those interested please address as above.

Miss Lydia Prichett Borden, Manoa, Pa., under date Jan. 11, 1907, writes that she found *Mnium spinulosum* B. & S. in full fruit and many plants with the spores apparently perfectly ripe on January 1st, this year. A friend also found *M. sylvaticum* in fruit and a few other species which usually fruit in spring. These were all found in Delaware County within a few miles of Philadelphia.

Since March 1st the following persons have qualified as Chapter members: No. 174. Dr. Marshall A. Howe, New York Botanical Garden, Bronx Park, New York City. No. 175. Mrs. L. E. C. Rohrabacher, Hotel Stander, Seattle, Wash. No. 176. Rev. C. H. Waddell, The Vicarage, Saintfield, Co. Down, Ireland. No. 177. Dr. I. Hagen, Natural History Museum, Trondhjem, Norway. No. 178. Dr. V. F. Brotherus, Helsingfors, Finland, Sweden.

In opening a new set of books it has been found advisable to send out all

unpaid accounts both to Chapter members and subscribers. This is not to be taken as a regular order of procedure, as the understanding is that all bills become due on the first of each January, and are to be paid without individual notice. Receipts are, however, always returned.

OFFERINGS.

(To Chapter Members only. For postage.)

- Rev. H. Dupret, Seminary of Philosophy, Montreal, Canada. *Dicranum fulvum* Hook.; *Paludella squarrosa* Brid., from near Montreal.
- Dr. J. F. Brenckle, Kulm, North Dakota. *Tortula mucronifolia*, B. & S., *Pterygoneurum cavifolium* (Ehrh.) Jur.
- Prof. Thomas A. Bonser, E. 12 Nora Avenue, Spokane, Wash. *Hypnum uncinatum* var. *symmetricum* R. & C. From Spokane.
- Dr. J. W. Bailey, 4541 Fourteenth Avenue, Seattle, Wash. *Dicranoweisia cirrhata* Lindb. From western Washington.
- Mr. E. B. Chamberlain, 35 West 59th street, New York City. *Plagiochila asplenoides* (L.) Dum.; *Preissia quadrata* (Scop.) Nees. Collected Pleasant Ridge, Maine.

AMERICAN HEPATICAE

When Prof. L. M. Underwood discontinued the issue of the Hepaticae Americanae he had a good deal of material on hand originally intended for future issues; he has very kindly turned this over to me and I am working it up, with Prof. A. W. Evans' valuable assistance in verifying and determining the species whenever necessary. This, with material of my own collecting, I propose to issue from year to year under the title of AMERICAN HEPATICÆ.

The first issue comprising decades I-II, is ready for distribution. Price, \$1.50. As only twenty sets of these decades are issued it will be necessary for subscribers to indicate their wishes at once.

Miss CAROLINE COVENTRY HAYNES ✻ HIGHLANDS, NEW JERSEY